




Background - ICU

■ Imperial College Model – ICUs overwhelmed



Innovation



Background - CPAP

- CPAP to improve outcomes
 - Qin Sun et al.
- Oxygen use


Lower mortality of COVID-19 by early recognition and intervention: experience from Jiangsu Province

Qin Sun¹, Haibo Qiu¹, Mao Huang^{2*} and Yi Yang^{1*}

BBC

Coronavirus doctor's diary: 'Our hospitals weren't made to use this much oxygen'

And Dr Tom Lawton, one of our ingenious intensive care consultants, ... Airway Pressure (CPAP) but some of the hospital's CPAP machines ...
7 Apr 2020




Physiotherapists

- Building “Team CPAP”
- Training and Support
- Managing Equipment




Outcomes



- 2x as much CPAP
- Nobody who could benefit was denied it
- Less than ½ the use of ICU & ventilators

- ICU not overwhelmed!

Outcomes



- Lower mortality despite a sicker population
- Successful transfer of Jiangsu's promise into an NHS model of care

After Wave 1

- Promotion around the NHS
 - Contact from other hospitals around the world!
 - BBC & other press interest
 - Research paper read over 6,000 times



The BMJ
Cite this as: *BMJ* 2021;374:n1950
<http://dx.doi.org/10.1136/bmj.n1950>
Published: 4 August 2021

Covid-19: CPAP reduces need for invasive mechanical ventilation in patients requiring oxygen, study finds

Elisabeth Mahase

Continuous positive airway pressure (CPAP) reduced the need for invasive mechanical ventilation in adults admitted to hospital with acute respiratory failure due to covid-19, a trial has found.¹

He said that the findings may be beneficial in systems with limited resources, such as oxygen.

The researchers noted a number of limitations to the trial, including that they did not achieve the planned

	Bradford patients	ISARIC patients	P
NIV use	29.5% (165 of 559)	15.9% (2670 of 16,805)	<0.0001
Critical care admission	7.2% (40 of 559)	16.5% (3001 of 18,183)	<0.0001
Intubation	5.0% (28 of 559)	9.8% (1658 of 16,866)	<0.0001
Hospital mortality	32.7% (183 of 559)	38.6% (5165 of 13,364)	0.005
Mortality for critical care patients	50% (20 of 40)	53.7% (958 of 1784)	NS
Median length of stay (IQR)	7 (3–14)	7 (unreported IQR) ³⁶	NS