



# Plenvu Bowel Preparation


Each box contains 3 sachets, making up dose 1 (1 sachet), and dose 2 (sachet A & B)

## Why do I need to take it?

Taking Plenvu before your appointment will give you a clean bowel.

## What to Expect

After you drink Plenvu you will have lots of watery bowel movements, so stay near a toilet because you may need to use it urgently.



## Reporting of Side Effects

### United Kingdom

If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in the package leaflet. You can also report side effects directly via the Yellow Card Scheme at [www.mhra.gov.uk/yellowcard](http://www.mhra.gov.uk/yellowcard). By reporting side effects you can help provide more information on the safety of this medicine.

A patient support item from Norgine.

Norgine Pharmaceutical Ltd, ARC Uxbridge, Building 01, Sanderson Road, Uxbridge UB8 1DH United Kingdom

E-mail: [ukisupport@norgine.com](mailto:ukisupport@norgine.com)

Tel: +44 (0)1895 453 681

Plenvu preparation leaflet adapted with kind permission from Norgine Pharmaceuticals LTD.

# Importance of Proper Cleansing

- A clean bowel will allow for a clear view of its inner lining which is important for both accurate diagnosis and treatment.
- Abnormal changes may be tiny, so even the smallest amount of faeces could potentially disguise something important.
- **If the bowel is not empty and clean enough, the procedure may have to be cancelled and repeated on another occasion at a later date.**

## How to Prepare Plenvu

### Dose 1



1 Empty dose 1 sachet in to a 500ml jug



2 Add water to make up 500ml and stir until fully dissolved



3 Sip very slowly over 60 minutes

### Dose 2




1 Empty dose 2 sachets (A & B) in to a 500ml jug



2 Add water to make up 500ml and stir until fully dissolved



3 Sip very slowly over 60 minutes




drink a minimum of **500ml** clear fluids with each dose. Alternating between this and the Plenvu solution is acceptable

# Patient Advice Clear Fluids

You must drink a minimum of 500ml extra clear fluids with each dose. Continue drinking clear fluids to keep yourself hydrated until 2 hours before your procedure and then stop drinking, clear drinks may include:




Water



Clear soups (no bits)



Tea & Coffee (without milk)



Soft drinks (cordial - not blackcurrant)

••••• Do Not Drink Alcohol •••••

## Plenvu may be easier to take when:

- Adding cordial to the additional clear fluids (to alter the flavour, but not blackcurrant as it will stain the bowel)
- It has been chilled once it has been made up

## Further advice for patients:

- Using wet wipes after each bowel movement
- Use a barrier cream to prevent a 'sore bottom' for example: petroleum jelly

## Hints & Tips



Sip the Plenvu solution very slowly over 60 minutes alongside a minimum of 500ml clear fluids (you can alternate between sips of Plenvu and clear fluid if you'd like)

Once dissolved (this may take up to 8 minutes) the solution can be refrigerated



Try using a straw



Stay hydrated with plenty of clear fluids